arXiv:math/0501204v1 [math.AG] 13 Jan 2005

A smooth, complete threefold with no nontrivial nef line bundles

Sam Payne

February 1, 2008

Abstract

We present a smooth, complete toric threefold with no nontrivial nef line bundles. This is a counterexample to a recent conjecture of Fujino.

Let X be a complete algebraic variety. A line bundle L on X is said to be nef if $L \cdot C$, the degree of L restricted to C, is nonnegative for every curve $C \subset X$. If X has no nontrivial nef line bundles, then X admits no nonconstant maps to projective varieties. Indeed, if $f : X \to \mathbb{P}^n$ is a nonconstant morphism, then $f^*\mathcal{O}(1)$ is a nef line bundle that is not isomorphic to $\mathcal{O}_X$. One example of a complete variety with no nontrivial nef line bundles, due to Fulton, is the singular toric threefold corresponding to the fan over a cube with one ray displaced, which has no nontrivial line bundles at all [Ful, pp.25-25, 72]. The literature also contains a number of examples of Moishezon spaces with no nontrivial nef line bundles. See [Nak, 3.3], [Kol, 5.3.14], [Ogu], and [Bon]. Recently, Fujino has given examples of complete, singular toric varieties with arbitrary Picard number that have no nontrivial nef line bundles. Fujino conjectured that $NE(X)$ is properly contained in $N_1(X)$ for any smooth, complete toric variety X. This conjecture is equivalent to the statement that every smooth, complete toric variety has a nontrivial nef line bundle [Fuj, Remark 3.1, Conjecture 4.5]. The purpose of this note is to give a counterexample to Fujino's conjecture; we present a smooth, complete toric threefold with no nontrivial nef line bundles.

Let Σ be the fan in $\mathbb{R}^3$ whose rays are generated by

$$\begin{array}{llll} v_1 = (1,1,1), & v_2 = (-1,1,1), & v_3 = (-1,-1,1), & v_4 = (1,-1,1), \\ v_5 = (1,0,-1), & v_6 = (0,1,-1), & v_7 = (-1,0,-1), & v_8 = (0,-1,-1), \end{array}$$

and whose maximal cones are $\langle v_1, v_2, v_3, v_4 \rangle$, $\langle v_5, v_6, v_7, v_8 \rangle$ and

$$\begin{array}{llll} \langle v_1, v_2, v_5 \rangle, & \langle v_2, v_3, v_6 \rangle, & \langle v_3, v_4, v_7 \rangle, & \langle v_4, v_1, v_8 \rangle, \\ \langle v_2, v_5, v_6 \rangle, & \langle v_3, v_6, v_7 \rangle, & \langle v_4, v_7, v_8 \rangle, & \langle v_1, v_8, v_5 \rangle. \end{array}$$

The nonzero cones of Σ are the cones over the faces of the following nonconvex polyhedron, which has a rotational symmetry about the vertical axis of order 4.



Let Δ be the fan obtained from Σ by successive stellar subdivisions along the rays spanned by $v_9 = (0,0,-1)$, $v_{10} = (0,0,1)$, and

$$v_{11} = (1,0,1), \quad v_{12} = (0,1,1), \quad v_{13} = (-1,0,1), \quad v_{14} = (0,-1,1).$$

We claim that $X = X(\Delta)$, the toric threefold corresponding to the fan Δ with respect to the lattice $\mathbb{Z}^3 \subset \mathbb{R}^3$, is smooth and complete and that X has no nontrivial nef line bundles. It is easy to check that $|\Delta| = \mathbb{R}^3$, so X is complete, and that each of the 24 maximal cones of Δ is spanned by a basis for $\mathbb{Z}^3$, so X is smooth. It remains to show that X has no nontrivial nef line bundles.

Suppose $D = \sum d_i D_i$ is a nef Cartier divisor, where D_i is the prime T-invariant divisor corresponding to the ray spanned by v_i. Since nef line bundles on toric varieties are globally generated, $\mathcal{O}(D)$ has nonzero global sections. Therefore we may assume that D is effective, i.e. each $d_i \geq 0$. We will show that $D = 0$. Note that $\langle v_2, v_5, v_6 \rangle$ is a cone in Δ, and

$$v_1 \;=\; 3v_2 + 4v_5 - 2v_6.$$

It then follows from the convexity of the piecewise linear function Ψ_D associated to D that

$$d_1 + 2d_6 \quad \geq \quad 3d_2 + 4d_5.$$

Similarly,

$$\begin{aligned} d_2 + 2d_7 &\quad \geq \quad 3d_3 + 4d_6, \\ d_3 + 2d_8 &\quad \geq \quad 3d_4 + 4d_7, \\ d_4 + 2d_5 &\quad \geq \quad 3d_1 + 4d_8. \end{aligned}$$

Adding the four above inequalities, we have

$$\begin{aligned} & d_1 + d_2 + d_3 + d_4 + 2(d_5 + d_6 + d_7 + d_8) \;\geq \\ & \qquad 3(d_1 + d_2 + d_3 + d_4) + 4(d_5 + d_6 + d_7 + d_8). \end{aligned}$$

Since all of the d_i are nonnegative, it follows that $d_i = 0$ for $1 \leq i \leq 8$. Since $v_1, \ldots, v_8$ positively span $\mathbb{R}^3$, we have $P_D = \{0\}$, and hence $D = 0$.

The above argument shows that any complete fan in $\mathbb{R}^3$ containing the cones $\langle v_2, v_5, v_6 \rangle$, $\langle v_3, v_6, v_7 \rangle$, $\langle v_4, v_7, v_8 \rangle$, and $\langle v_1, v_5, v_8 \rangle$ corresponds to a complete toric variety with no nef line bundles. In particular, stellar subdivisions of the remaining cones in Δ lead to examples of smooth, complete toric threefolds with arbitrarily large Picard number and no nontrivial nef line bundles.

References

[Bon] Sur des variétés de Moishezon dont le groupe de Picard est de rang un, Bull. Soc. math. France, **124** 503–521.

[Fuj] O. Fujino, On the Kleiman-Mori cone, preprint 2005. math.AG/0501055

[Ful] W. Fulton, *Introduction to Toric Varieties*, Annals of Math. Studies **131**, Princeton University Press 1993.

[Kol] J. Kollár, Flips, flops, minimal models, etc., Surveys in Diff. Geom. **1** (1991) 113–199.

[Nak] I. Nakamura, Moishezon threefolds homeomorphic to $\mathbb{P}^3$, J. Math. Soc. Japan **39** (1987) 521–535.

[Ogu] K. Oguiso, Two remarks on Calabi-Yau Moishezon threefolds, J. Reine Angew. Math. **452** (1994) 153–161.